arXiv:cond-mat/9809129v1 [cond-mat.str-el] 8 Sep 1998

Subject classification: 71.55.-i; 75.50.Ee; S10.15

Ferron-like states in $YBa_2Cu_3O_{6+x}$

P. Rubin and A. Sherman

Institute of Physics, University of Tartu, Riia 142,
EE-2400 Tartu, Estonia

With the use of the Hubbard model bound hole states in $YBa_2Cu_3O_{6+x}$ are studied. For the parameters of this crystal the exchange interaction between the spin-carrying chain ion O^- and $Cu-O$ plane sites is shown to ensure the formation of a large ferromagnetically ordered clusters around holes in the plane.

1. Introduction

Over the past decade the magnetic structure of cuprate perovskites has been studying very intensively in the hope that it might provide insight in the physical origin of high-temperature superconductivity. It is presently well established that CuO_2 planes of undoped and lightly doped cuprates are antiferromagnetically ordered. Some experiments in such crystals, for example magnetization measurements in $YBa_2Cu_3O_{6+x}$ [1], suggest the existence of local ferromagnetically ordered regions around holes on the antiferromagnetic background of CuO_2 planes. In application to another type of crystals such excitations were first considered in Ref. 2 and were called ferrons. Ferron models of high-T_c materials were discussed in Refs. 3,4. The explanation of the ferron formation, suggested in Ref. 1, is based on the comparison of antiferromagnetic and ferromagnetic interactions created by a hole localized on an oxygen orbital. This idea goes back to work [5] where it was used for the explanation of the observed destruction of the long-range antiferromagnetic order at small hole concentrations. The mechanism suggested in the latter work is essentially based on the supposition that nonhybridized oxygen and copper orbitals are appropriate starting states for the consideration. However as shown in Ref. 6, low-lying states are strongly hybridized. As a consequence, the mechanism of Ref. 5 is not appropriate for explaining the destruction of the long-range ordering. For similar reasons this mechanism may not be responsible for the ferron formation.

Notice that the mentioned strong hybridization of oxygen and copper orbitals and the crystal structure of CuO_2 planes open the way to reduce the Emery Hamiltonian for these planes to the more simple t-J or Hubbard Hamiltonians with only one effective orbital per plane sell [6]. In the t-J model the ferron formation was considered both by the exact diagonalization of small clusters [7, 8, 9] and in an infinite plane with the use of the spin-wave approximation [10, 11]. Both approaches showed that in the periodic lattice ferrons are energetically unfavorable for parameters of cuprate perovskites. The next step is to check whether point defect can stabilize ferrons. In our recent paper [12] we have considered local states generated by the interstitial O^{2-} ion in $La_2CuO_{4+\delta}$ and found that this type of defects is unable to stabilize ferrons. In $YBa_2Cu_3O_{6+x}$, where ferrons were experimentally observed, for small x oxygen ions start to fill chain sites and can be considered as interstitial defects above the CuO_2 plane (see Fig. 1). Their valency is equal to -2 or -1 [13] and in the latter case the defect has the spin 1/2. It can be expected that, in contrast to the spinless O^{2-} ion, such type of defects can stabilize states with a large spin, i.e. ferrons, due to an arising exchange interaction.

2. Basic Formulae

To check this idea we use the one-band Hubbard model for the description of a CuO_2 plane. The model Hamiltonian reads

$$H_0 = U \sum_{\mathbf{l}} n_{\mathbf{l}\uparrow} n_{\mathbf{l}\downarrow} - t \sum_{\mathbf{l}\mathbf{a}\sigma} c^\dagger_{\mathbf{l}\sigma} c_{\mathbf{l}+\mathbf{a}\sigma}, \qquad (1)$$

where the operator $c^\dagger_{\mathbf{l}\sigma}$ creates an electron in the effective orbital on the plane site $\mathbf{l}$ (the orbitals are centered at copper sites) with the spin σ, $\mathbf{a} = (0, \pm a), (\pm a, 0)$, where a is the in-plane copper distance, $n_{\mathbf{l}\sigma} = c^\dagger_{\mathbf{l}\sigma} c_{\mathbf{l}\sigma}$, U and

t are the Coulomb repulsion and the hopping matrix element, respectively. A chain oxygen O$^-$ is situated at O1 position and we add the following terms to the Hamiltonian

$$H_1 = \Delta \sum_{\sigma} a_{\sigma}^{\dagger} a_{\sigma} - \sum_{\mathbf{m}\sigma} t_{\mathbf{m}} \left(a_{\sigma}^{\dagger} c_{\mathbf{m}\sigma} + c_{\mathbf{m}\sigma}^{\dagger} a_{\sigma} \right) + U_p a_{\uparrow}^{\dagger} a_{\uparrow} a_{\downarrow}^{\dagger} a_{\downarrow}, \tag{2}$$

where $a_{\sigma}^{\dagger}$ is the p-electron creation operator in the O$^-$ chain site, Δ is the difference in chain and plane site energies, U_p is the Coulomb repulsion on the chain oxygen, and $t_{\mathbf{m}}$ is the hopping matrix element between chain and plane sites. Summation over $\mathbf{m}$ proceeds over the plane sites in the vicinity of the "defect" chain oxygen site (these plane sites are shown in Fig. 1) and we suppose that $t_{\mathbf{m}} \approx t$ for these sites.

For YBa$_2$Cu$_3$O$_{6+x}$ the parameters in Eqs. (1) and (2) can be estimated as $t_{\mathbf{m}} \approx t \approx 1\,\mathrm{eV}$, $\Delta \approx -2\,\mathrm{eV}$, $U \approx 10\,\mathrm{eV}, U_p \approx 4.5\,\mathrm{eV}$ [10, 14]. For such parameters the correlation terms in Eqs. (1),(2) and the first term in Eq. (2) can be used as the unperturbed Hamiltonian and the second, hopping terms, as the perturbation in the perturbation theory expansion. The resulting effective Hamiltonian reads

$$H_{\text{eff}} = P \left(-t \sum_{\mathbf{la}\sigma} c_{\mathbf{l}\sigma}^{\dagger} c_{\mathbf{l+a}\sigma} + \frac{J}{2} \sum_{\mathbf{la}} \mathbf{S}_{\mathbf{l+a}} \mathbf{S}_{\mathbf{l}} + \sum_{\mathbf{m}} J_{\mathbf{m}} \mathbf{S}_{\mathbf{m}} \mathbf{S}_{\mathbf{o}} \right) P,$$

$$J = \frac{4t^2}{U}, \qquad J_{\mathbf{m}} = \frac{2t_{\mathbf{m}}^2 (U + U_p)}{(\Delta + U_p)(U - \Delta)}, \tag{3}$$

where

$$S_{\mathbf{l}}^z = \sum_{\sigma} \sigma c_{\mathbf{l}\sigma}^{\dagger} c_{\mathbf{l}\sigma}/2, \quad S_{\mathbf{l}}^{\sigma} = S_{\mathbf{l}}^x + i\sigma S_{\mathbf{l}}^y = c_{\mathbf{l}\sigma}^{\dagger} c_{\mathbf{l}-\sigma},$$

are the components of the spin operators $\mathbf{S}_{\mathbf{l}}$ on the plane sites. The analogous Hamiltonian was derived in [15]. The components of the spin operator $\mathbf{S}_{\mathbf{o}}$ on the chain site are defined analogously through the operators a_{σ}. P is the projection operator on the subspace where the chain level is singly occupied and the plane levels are singly occupied or empty. The first two terms in Eq. (3) represent the one-band t-J Hamiltonian on a square lattice. The summation over $\mathbf{l}$ proceeds over all plane sites, while the summation over $\mathbf{m}$, over the "defect" plane sites. In Eq. (3), the "defect" term can be represented as the sum of terms with S^z and $S^{\pm}$ spin components. The term with S^z components plays the main role in the ferron stabilization, whereas the terms with $S^{\pm}$ components lead only to additional lowering of energy levels. For this reason for simplicity we take into account only components with S^z in the "defect" term in Eq. (3). Let us suppose that the O1 site is occupied by an electron with $S^z = 1/2$. In this case the effective Hamiltonian can be rewritten in the form

$$H_{\text{eff}} = -t \sum_{\mathbf{la}\sigma} c^{\dagger}_{\mathbf{l}\sigma} c_{\mathbf{l+a}\sigma} + \frac{J}{2} \sum_{\mathbf{l},\mathbf{a}} \mathbf{S}_{\mathbf{l+a}} \mathbf{S}_{\mathbf{l}} + \frac{1}{2} \sum_{\mathbf{m}} J_{\mathbf{m}} S^z_{\mathbf{m}}. \tag{4}$$

Since the sign of $J_{\mathbf{m}}$ is the same for all sites in the "defect" region, the last term can stabilize ferrons.

We use the spin-wave approximation [10, 16] for the description of spin excitations in the plane. The approximation reduces to neglecting terms of the third and higher orders in the spin-wave operators $b_{\mathbf{m}}$ introduced into Eq. (4) by the Holstein-Primakoff formulas

$$S^{+1}_{\mathbf{l}} = \Phi_{\mathbf{l}} b_{\mathbf{l}} P^{+1}_{\mathbf{l}} + b^{\dagger}_{\mathbf{l}} \Phi_{\mathbf{l}} P^{-1}_{\mathbf{l}}, \;\; S^{-1}_{\mathbf{l}} = (S^{+1}_{\mathbf{l}})^{\dagger}, \;\; S^z_{\mathbf{l}} = e^{i\mathbf{\Pi l}} \left(\frac{\sum_{\sigma} n_{\mathbf{l}\sigma}}{2} - b^{\dagger}_{\mathbf{l}} b_{\mathbf{l}} \right),$$

where $\Phi_{\mathbf{l}} = \left(1 - b^{\dagger}_{\mathbf{l}} b_{\mathbf{l}}\right)^{1/2}$, $\mathbf{\Pi} = (\pi/a, \pi/a)$, and $P^{\sigma}_{\mathbf{l}} = [1 + \sigma \exp(i\mathbf{\Pi l})]/2$.

Let us introduce the hole creation operator $h^{\dagger}_{\mathbf{l}} = \sum_{\sigma} P^{\sigma}_{\mathbf{l}} c_{\mathbf{l}\sigma}$ for the Néel state $|\mathcal{N}\rangle$. This classical Néel state can be determined by the condition $b_{\mathbf{l}}|\mathcal{N}\rangle = 0$ (for the second Néel state analogous formulas are obtained by substituting $P^{\sigma}_{\mathbf{l}}$ with $1 - P^{\sigma}_{\mathbf{l}}$). After the unitary transformation which diagonalizes the Heisenberg part of the Hamiltonian [10] (the second term in Eq. (4)) the effective Hamiltonian acquires the form

$$\begin{aligned} H_{\text{eff}} &= \sum_{\mathbf{ll'a}} t \left[h_{\mathbf{l+a}} h^{\dagger}_{\mathbf{l}} b^{\dagger}_{\mathbf{l-l'}} (u_{\mathbf{l'+a}} + v_{\mathbf{l'}}) + \text{H.c.} \right] + \frac{J}{2} \sum_{\mathbf{ll'}} \omega_{\mathbf{l'}} b^{\dagger}_{\mathbf{l}} b_{\mathbf{l+l'}} \\ &- \sum_{\mathbf{m}} \left(J_{\mathbf{1m}}\, \tilde{n}_{\mathbf{m}} + J_{\mathbf{2m}}\, b^{\dagger}_{\mathbf{m}} b_{\mathbf{m}} \right) \exp(i\mathbf{\Pi m}), \end{aligned} \tag{5}$$

where $u_{\mathbf{l}}$, $v_{\mathbf{l}}$ and $\omega_{\mathbf{l}}$ are the Fourier transforms of $\cosh(\alpha_{\mathbf{k}})$, $-\sinh(\alpha_{\mathbf{k}})$, and $\omega_{\mathbf{k}} = 4\sqrt{1 - \gamma^2_{\mathbf{k}}}$, respectively, $\alpha_{\mathbf{k}} = (1/8)\, \ln\left[(1 + \gamma_{\mathbf{k}})/(1 - \gamma_{\mathbf{k}})\right]$, $\gamma_{\mathbf{k}} = [\cos(k_x a) + \cos(k_y a)]/2$, $\tilde{n}_{\mathbf{m}} = h^{\dagger}_{\mathbf{m}} h_{\mathbf{m}}$. Due to the fast decrease of $u_{\mathbf{l'+a}} + v_{\mathbf{l'}}$ and $\omega_{\mathbf{l'}}$ with the growth of $|\mathbf{l'}|$ only the components with $\mathbf{l'} = (\pm a, 0), (0, \pm a)$ for the sum in the first, kinetic energy term and the components with $\mathbf{l'} = (0,0), (\pm a, \pm a), (\pm 2a, 0), (0, \pm 2a)$ for $\omega_{\mathbf{l'}}$ in the second, magnetic energy term are retained in Eq.(5) in the subsequent calculations. In this equation, $J_{\mathbf{1m}} = J_{\mathbf{m}}/4$, $J_{\mathbf{2m}} = 2(u_{\mathbf{0}}^2 + v_{\mathbf{0}}^2)J_{\mathbf{1m}}$ where $u_{\mathbf{0}}^2 + v_{\mathbf{0}}^2 \approx 1.16$.

To investigate the lowest eigenvalues and eigenfunctions of Hamiltonian (5) we use the modified Lanczos algorithm [10, 17]. In one step of this algorithm we determine a final state $|\mathbf{f}\rangle$ from an initial state $|\mathbf{i}\rangle$ according to the procedure

$$\langle\mathbf{i}|\mathbf{i}\rangle = 1, \;\; \mathcal{E}_i = \langle\mathbf{i}|\mathcal{H}|\mathbf{i}\rangle, \;\; \mathcal{V}|\mathbf{f}\rangle = (\mathcal{H} - \mathcal{E}_i)|\mathbf{i}\rangle. \tag{6}$$

Here $\mathcal{V}$ is the normalization constant which is determined from the condition $\langle\mathbf{f}|\mathbf{f}\rangle = 1$. After one Lanczos step the energy is minimized in the subspace of the states $|\mathbf{i}\rangle$ and $|\mathbf{f}\rangle$ (it follows from (6) that $\langle\mathbf{i}|\mathbf{f}\rangle = 0$). The obtained function $|\mathbf{i}'\rangle = c_1|\mathbf{i}\rangle + c_2|\mathbf{f}\rangle$ which minimizes the energy is used as the initial function in the next Lanczos step. This procedure is continued until the necessary accuracy of the eigenvalue is reached. If the initial function $|\mathbf{i}\rangle$ is characterized by some spin projection S^z, $|\mathbf{f}\rangle$ also corresponds to the same S^z. Therefore, it is possible to obtain the lowest eigenvalues of Hamiltonian (8) with a given S^z by starting from a state with the respective S^z. Notice that states with $S^z > 1/2$ contain flipped spins near the hole and correspond to ferrons.

The number of spin configurations which is necessary for describing the states $|\mathbf{i}\rangle$ and $|\mathbf{f}\rangle$ grows from step to step. To avoid the overflow of the computer storage and to reduce the computation time, in each Lanczos step the state $|\mathbf{i}'\rangle$ is restricted to the j configurations with the largest amplitudes [10]. After normalization this restricted state is used as the initial state for the next recursion step of the Lanczos procedure (6). We used $j = 200$. The $|\mathbf{f}\rangle$ state obtained from such $|\mathbf{i}'\rangle$ state and used for the calculation of the eigenvalue contains of the order of 10^4 spin configurations. This is comparable to the number of configurations used in exact diagonalization studies of small lattices. To check the accuracy of the obtained results in some cases we increased j up to 250 which led only to small quantitative changes (of the order of $10^{-4}t$) in the eigenvalue.

3. Results and Discussion

Using the above-described algorithm we have calculated the lowest one-hole states characterized by $S^z \leq 13/2$. For parameters of cuprate perovskites the ratio J/t can be estimated to lie in the range 0.2 - 0.5 [10]. According to Eq. (3) the value of $J_{\mathbf{m}}/t$ for nearest neighbours of O1 oxygen site can be estimated to be equal 1. For other sites in the "defect" region the values of $J_{\mathbf{m}}/t = 0.7$ is accepted. The results of our calculations for the lowest one-hole eigenvalues are shown in Fig. 2. As seen, the interaction between the O^- ion spin with plane spins leads to the ferron formation: the lowest state has $S^z = 7/2$. The optimal ferron size can be expected to grow for two and larger number of adjacent O^- ions in a chain.

In summary, with the use of the spin-wave approximation and the Lanczos algorithm the localized hole states induced by an O^- chain ion in $YBa_2Cu_3O_{6+x}$ were investigated. The lowest hole state is characterized by the z component of the total spin $S^z = 7/2$ which corresponds to a comparatively large ferromagnetic region with 3 overturned spins around a hole. This means that the spin-carrying O^- ion can stabilize ferrons.

Acknowledgements This work was supported by the Estonian Science Foundation under Grant ETF-2688.

References

[1] S. L. Gnatchenko, A. M. Ratner, M. Baran, R. Szymczak, and H. Szymczak, Phys. Rev. B **55**, 3876 (1997).

[2] P. G. de Gennes, J. Phys. Radium **23**, 630 (1962).

[3] V. L. Pokrovskii, G. V. Uimin, and D. V. Khveshchenko, JETP Lett. **46 Suppl.**, S113 (1987).

[4] V. Hizhnyakov and E. Sigmund, Physica C **156**, 655 (1988).

[5] R. J. Birgeneau, M. A. Kastner, and A. Aharony, Z. Phys. B **71**, 57 (1988).

[6] F. C. Zhang and T. M. Rice, Phys. Rev. B **37**, 3759 (1988).

[7] J. Bonča, P. Prelovšek, and I. Sega, Phys. Rev. B **39**, 7074 (1989).

[8] E. Dagotto, A. Moreo, and T. Barnes, Phys. Rev. B **40**, 6721 (1989).

[9] Y. Hasegawa and D. Poilblank, Phys. Rev. B **40**, 9035 (1989).

[10] A. V. Sherman, Solid State Commun. **76**, 321 (1990), Physica C **171** 395 (1990).

[11] J. Sabczynski, M. Schreiber, and A. Sherman, Phys. Rev. B **48**, 543 (1993).

[12] P. Rubin, A. Sherman, and M. Schreiber, Phys. Rev. B **57**, 10299 (1998).

[13] D. R. Grempel, H. Haugerud, G. Uimin, and W. Selke, Czechoslovak Journal of Physics **46 Suppl. S2**, 967 (1996).

[14] A. K. McMahan, J. F. Annett, and R. M. Martin, Phys. Rev. B **42**, 6268 (1990).

[15] T. Schork and P. Fulde, Phys. Rev. B **50**, 1345 (1994).

[16] S. Schmitt-Rink, C. M. Varma, and A. E. Ruckenstein, Phys. Rev. Lett. **60**, 2793 (1988).

[17] E. Dagotto and A. Moreo, Phys. Rev. D **31**, 865 (1985).

Figure captions.

Fig. 1. The “defect” region in the vicinity of the chain O$^-$ ion (O1).

Fig. 2. The dependence of energies of the localized hole states induced by the O$^-$ ion on S^z.


O1


-2.4
-2.8
-3.2
-3.6
E/t
0.0
2.0
4.0
6.0
8.0
S_z